

## CLINICAL RESEARCH

## Interventional Cardiology

# Cause of Death Within 30 Days of Percutaneous Coronary Intervention in an Era of Mandatory Outcome Reporting

Bhuvnesh Aggarwal, MD,\* Stephen G. Ellis, MD,† A. Michael Lincoff, MD,† Samir R. Kapadia, MD,† Joseph Cacchione, MD,† Russell E. Raymond, DO,† Leslie Cho, MD,† Christopher Bajzer, MD,† Ravi Nair, MD,† Irving Franco, MD,† Conrad Simpfordorfer, MD,† E. Murat Tuzcu, MD,† Patrick L. Whitlow, MD,† Mehdi H. Shishehbor, DO, MPH†

*Cleveland, Ohio*

## Objectives

This study sought to ascertain causes of death and the incidence of percutaneous coronary intervention (PCI)-related mortality within 30 days.

## Background

Public reporting of 30-day mortality after PCI without clearly identifying the cause may result in operator risk avoidance and affect hospital reputation and reimbursements. Death certificates, utilized by previous reports, have poor correlation with actual cause of death and may be inadequate for public reporting.

## Methods

All patients who died within 30 days of a PCI from January 2009 to April 2011 at a tertiary care center were included. Causes of death were identified through detailed chart review using Academic Research Consortium consensus guidelines and compared with reported death certificates. The causes of death were divided into cardiac and noncardiac and PCI and non-PCI-related categories.

## Results

Of the 4,078 PCI, 81 deaths (2%) occurred within 30 days. Of these, 58% died of cardiac and 42% of noncardiac causes. However, only 42% of 30-day deaths were attributed to PCI-related complications. Patients with non-PCI-related, compared with PCI-related, death presented with a higher incidence of cardiogenic shock (15 of 47 [32%] vs. 2 of 34 [6%];  $p < 0.01$ ) and cardiac arrest (19 of 47 [40%] vs. 1 of 34 [3%];  $p < 0.01$ ). Death certificates had only 58% accuracy (95% confidence interval: 45% to 72%) for classifying patients as experiencing cardiac versus noncardiac death.

## Conclusions

Less than one-half of 30-day deaths are attributed to a PCI-related complication. Death certificates are inaccurate and do not report PCI-related deaths, which may represent a better marker of PCI quality. (J Am Coll Cardiol 2013;62:409–15) © 2013 by the American College of Cardiology Foundation

Public reporting of outcomes after percutaneous coronary intervention (PCI) is likely to be made standard practice as advocated by the Centers for Medicare and Medicaid Services (1). Implementation of this policy will lead to several important consequences. First, Medicare and Medicaid reimbursements for healthcare professionals and institutions will be based on their outcomes (2). Second, mandatory

reporting and disclosure of these data will affect the credibility and reputation of a healthcare provider, hopefully helping patients to make informed decisions about their healthcare choices (3). Finally, analysis of the outcomes data will be useful in determining appropriateness and hence cost-effectiveness of performing these procedures (4).

See page 416

From the \*Department of Internal Medicine, Medicine Institute, Cleveland Clinic, Cleveland, Ohio; and the †Department of Cardiovascular Medicine, Heart and Vascular Institute, Cleveland Clinic, Cleveland, Ohio. Dr. Cacchione has served as a member of the advisory boards of Humana and UnitedHealthcare. Dr. Nair has received speaker's fees, research support, and honoraria from Boston Scientific, Abbott Vascular, and St. Jude Medical. Dr. Shishehbor has served as an uncompensated consultant and educator for Medtronic, Abbott Vascular, BARD, Bayer HealthCare, and Spectranetics. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose.

Manuscript received December 12, 2012; revised manuscript received February 15, 2013, accepted March 5, 2013.

Whereas publicly reported outcomes could be considered surrogate markers for healthcare quality, thus far, such disclosures have resulted in mixed reactions from healthcare providers (5–7). In New York and Massachusetts where PCI outcomes were first publicly reported, a strong selection bias toward avoidance of PCI in high-risk patients was subsequently described (8,9). Because of these potentially adverse consequences, many have advocated the importance

Abbreviations and Acronyms

ARC = Academic Research Consortium

CI = confidence interval

GI = gastrointestinal

PCI = percutaneous coronary intervention

STEMI = ST-segment elevation myocardial infarction

these risk scores were derived from mortality data obtained through death certificates, which are often inaccurate in precisely identifying the cause of death (15–17). Another limitation is lack of data on PCI versus non-PCI-related mortality, which may be a better measure of operator performance and PCI quality. For these reasons, we conducted a detailed chart review to identify the causes of 30-day post-PCI deaths employing standard definitions from globally recognized Academic Research Consortium (ARC) consensus guidelines (18,19).

Methods

**Study population.** All patients who underwent PCI from January 2009 to April 2011 at a single tertiary care center were identified through the institutional review board–approved institutional PCI registry. Baseline characteristics, cardiac history and risk factors, medications, other chronic medical illnesses, and angiographic and procedural data were prospectively obtained and recorded. Thirty-day deaths after PCI were identified after querying the Social Security Death Index. Circumstances surrounding death such as the decision to withdraw care and site of death were obtained. Medical charts were retrospectively reviewed for cause of death together by 2 clinicians (B.A. and M.H.S.) using standard ARC definitions to prevent any bias. Cause of death by chart review was compared with death certificates when available.

**Definitions.** Cardiogenic shock on presentation was defined as systolic blood pressure persistently <90 mm Hg or the need for inotropic support or intra-aortic balloon

pumping to maintain systolic blood pressure >90 mm Hg, in the presence of adequate left ventricular filling pressure and systemic hypoperfusion (18). Neurological dysfunction was defined as Glasgow Coma Scale score <9 or unresponsiveness to any stimuli with absence of brain stem reflexes (in cases where the Glasgow Coma Scale was not checked).

Cause of death was defined as the precipitating factor for the patient’s demise, independent of the presenting medical problem on admission (20). There were 19 deaths after initial discharge of whom 8 died after readmission. An attempt to obtain information regarding 11 out-of-hospital deaths was made through a nurse phone call to the patients’ families. Of these, no record could be obtained, and the families could not be reached for 5 patients.

The causes of death were divided into cardiac- and noncardiac-related and PCI- and non-PCI-related categories. Cardiac death was defined as any death due to proximate cardiac cause (such as myocardial infarction, low output failure, or fatal arrhythmia), unwitnessed death, death from unknown cause, and all procedure-related deaths, including those related to concomitant treatment. Noncardiac causes were divided into respiratory, infectious, neurological, gastrointestinal (GI), renal and hemato-oncological categories (Table 1). All deaths were considered cardiac unless an unequivocal noncardiac cause could be established. In cases without reasonable clinical evidence toward one or the other, death was determined noncardiac only if stable cardiac pump function was determined by pulmonary artery catheterization.

PCI-related death was defined as death from complication of procedure such as vascular dissection, aneurysm, perforation, bleeding, renal failure, and definite or probable stent thrombosis. Stent thrombosis was defined as “definite” after angiographic confirmation or “probable” in case of any unexplained death within 30 days or any myocardial infarction that was related to documented acute ischemia in the territory of the implanted stent (18). Any bleeding was considered PCI-related if it occurred within 72 h of the procedure, and death from such bleeding was labeled as a complication of PCI (21).

**Statistical analysis.** PCI and non-PCI-related deaths were compared with regard to demographics, past medical history, presenting history, indication for procedure, and the

Table 1 Definitions of Noncardiac Causes of Death	
Cause of Death	Definition
Infectious	Death from severe sepsis or septic shock as defined by the Society of Critical Care Medicine/American College of Chest Physicians consensus conference guidelines (19).
Neurological*	Death due to anoxic brain injury prior to PCI, cerebrovascular accident, or brain death from any cause.
Pulmonary	Death after worsening respiratory status due to primary lung pathology including acute respiratory distress syndrome.
Gastrointestinal	Death from massive gastrointestinal bleeding, as complication of liver disease, cancer, or gastrointestinal perforation not related to dual antiplatelet therapy.
Hemato-oncological	Death from life-threatening hemorrhage (except gastrointestinal and intracranial hemorrhage) or due to an advanced cancer, its complication, or withdrawal of care due to concerns regarding poor prognosis associated with cancer.
Renal†	Death due to complication of renal failure such as fluid overload, acidosis, and electrolyte disturbances.

\*In patients with cardiac arrest, death was adjudicated to be neurological only if neurological dysfunction was documented prior to the start of percutaneous coronary intervention (PCI). In addition, the patient was also required to have a successful PCI with restoration of stable cardiac pump function as determined by either a pulmonary artery catheterization or lack of inotropic support. In equivocal cases or if no such data were available, cause of death was considered to be cardiac. †Excluding patients with contrast-induced nephropathy from iodinated contrast administered during PCI.

site of death. Continuous variables were presented as mean  $\pm$  SD or median with interquartile range, and categorical variables were presented as percentage affected. Fisher exact test for categorical variables and Wilcoxon rank order test for continuous variables were used to compare differences in categories. Accuracy, sensitivity, and specificity of death certificates for classifying patients as experiencing cardiac versus noncardiac death were calculated after assuming chart review diagnosis as the gold standard. All tests were 2-tailed and a  $p$  value  $<0.05$  was considered significant.

## Results

A total of 4,078 PCI (including emergent, urgent, elective, and salvage) were performed from January 2009 to April 2011. All-cause mortality within 30 days of PCI was 2% ( $n = 81$ ), and a cause of death could be established in 76 patients. Death records were not available for 5 patients who died after hospital discharge. Forty-seven patients (58%) died from cardiac and 34 (42%) from noncardiac causes (Fig. 1). Fifty-three patients presented with cardiogenic shock and underwent PCI, of these, 17 (32%) died within 30 days. Similarly, mortality for patients presenting with ST-segment elevation myocardial infarction (STEMI) and undergoing PCI was 7% (37 of 535 patients), whereas that for patients presenting with cardiac arrest was 41% (20 of 49 patients).


Of the noncardiac causes of death, complications of infection such as septic shock or severe sepsis were most common. About one-half of these had evidence of active infection prior to PCI (as demonstrated by a positive blood culture within 72 h of PCI). Death due to a neurological

cause was the next most prevalent noncardiac cause (Fig. 1). The majority of these were from withdrawal of care after anoxic brain injury, which had likely occurred prior to PCI as a result of cardiac arrest or shock.

Of the cardiac deaths, 34 patients died from PCI-related complications and 13 from non-PCI-related cardiac complications mainly related to pump dysfunction. The majority of PCI-related deaths were attributed to probable stent thrombosis (Fig. 2). Four patients died from fatal bleeding within 72 h of PCI with 2 cases of intracranial hemorrhage and 1 case each of massive GI bleeding and retroperitoneal hemorrhage. Lastly, 3 patients died from coronary dissection, and 2 died from complications of renal failure attributed to iodinated contrast administered with the procedure. Both patients had documented stable renal function prior to PCI.


Baseline and angiographic characteristics of PCI and non-PCI-related deaths (including noncardiac) are shown in Table 2. Both groups were comparable in terms of baseline demographics and pre-existing cardiac risk factors. However, patients who died from a non-PCI-related death presented more often with cardiogenic shock ( $p < 0.01$ ) or cardiac arrest ( $p < 0.01$ ) and had a higher prevalence of New York Heart Association functional class IV heart failure ( $p < 0.01$ ) prior to hospitalization (Table 2). In addition, non-PCI-related deaths occurred more often during index admission than did PCI-related deaths ( $p < 0.01$ ) (Fig. 3).

Death certificates were available in 53 patients. There was poor correlation between death certificates and chart review for classifying death into cardiac versus noncardiac death. When compared with chart review, death certificates had a sensitivity of 69% (95% confidence interval [CI]: 52% to



**Figure 1** Etiology of 30-Day Mortality After PCI Based on Chart Review

Etiology of death (in percentage) divided into percutaneous coronary intervention (PCI)-related, cardiac non-PCI-related, neurological, infection, pulmonary, gastrointestinal, and hemato-oncological causes.



86%), specificity of 46% (95% CI: 26% to 66%), and an accuracy of 58% (95% CI: 45% to 72%) (Fig. 4).

## Discussion

This is the first report in the past decade identifying etiology of death within the critical 30-day period after a PCI from a thorough chart review and using standard ARC guidelines. In our cohort, cardiac causes contributed to 58% of all deaths after PCI, and only 42% were attributed to complications associated with the procedure. Even this was perhaps an overestimation because all unexplained deaths within 30 days of PCI were considered probable stent thrombosis in this analysis. This is in contrast to previously reported data using death certificates where a much higher proportion of deaths were attributed to cardiac- and PCI-related complications (15,22–24). The highest-risk patients (cardiogenic shock and cardiac arrest) who may potentially experience risk avoidance rarely died from a PCI-related complication. In addition, a poor correlation between cause of death determined by chart review and that reported by death certificates was observed. These results highlight the limitations of 30-day mortality reporting when death certificates are used and emphasize the need for identifying PCI-related deaths rather than cardiac deaths alone.

Mortality is an outcome measure that is most likely to influence reimbursements and the public reputation of an institution. Therefore, it is important to classify causes of death into categories that truly reflect a physician and an institution's competence and acts as an indicator of performance. Most PCI and acute coronary syndrome outcome studies classify

deaths as from either cardiac or noncardiac causes rather than PCI- or non-PCI-related causes. Such classification may be misleading as institutional reporting of low cardiac death rates may be a result of risk avoidance (refusing PCI in high-risk patients such as those with STEMI, cardiogenic shock, or cardiac arrest) or a result of referral of such patients to different centers (6,7,25). We propose classifying post-PCI deaths as PCI- or non-PCI-related, which better represents quality of PCI and operator performance.

The overall mortality after PCI for a given provider is greatly influenced by the severity of a patient's illness and presentation. This is currently determined by several risk scores, but there remain concerns among interventionalists regarding the validity of these to accurately predict mortality after PCI (26). Because the expected in-hospital mortality after PCI is very low, this concern is especially relevant to the patients at the highest risk of death (27). Lack of trust in the current risk prediction models has led to avoidance of high-risk cases by clinicians in centers where outcome reporting was made mandatory. This was evident from the outcomes data published from the New York State PCI registry, which revealed that in-hospital mortality after PCI had declined from 0.90% in 1997 (28) to 0.58% in 2003 (29), a reduction of 36%. Although this might be superficially interpreted as due to a significant quality improvement, closer analysis revealed a simultaneous trend toward avoidance of PCI in patients presenting with cardiogenic shock (8). Similar results were observed in data from the Massachusetts PCI registry when outcome reporting became mandatory in that state (27). A "compassionate use" variable was proposed to account for

**Table 2** Patient Characteristics According to Etiology of Death by Chart Review

	PCI-Related Death (n = 34)	Non-PCI-Related Death (n = 47)	p Value
<b>Baseline characteristics and cardiac risk factors</b>			
Age, yrs	69 ± 14	71 ± 11	0.43
Male	19 (56)	28 (60)	0.82
Previous myocardial infarction	20 (59)	32 (68)	0.48
Previous percutaneous coronary intervention	14 (41)	15 (32)	0.48
Previous coronary artery bypass graft	3 (9)	9 (19)	0.23
Diabetes mellitus	14 (41)	24 (51)	0.50
History of renal failure	5 (15)	11 (23)	0.41
Current smoker	4 (12)	11 (23)	0.25
Peripheral vascular disease	8 (24)	9 (19)	0.78
<b>Cardiac characteristics at admission</b>			
Ejection fraction	39 ± 15	36 ± 15	0.58
STEMI	13 (38)	24 (51)	0.27
Non-STEMI	5 (15)	14 (30)	0.18
Unstable angina	5 (15)	2 (4)	0.12
Cardiogenic shock	2 (6)	15 (32)	<0.01
Cardiac arrest	1 (3)	19 (40)	<0.01
<b>Other comorbid conditions</b>			
History of stroke/transient ischemic attack	5 (15)	11 (23)	0.41
Neurological dysfunction	3 (9)	14 (30)	0.03
Infection pre-procedure	1 (3)	6 (13)	0.23
Dementia	4 (12)	0 (0)	0.03
Outside hospital transfer	20 (59)	25 (53)	0.66
<b>New York Heart Association functional class</b>			
I	1 (3)	3 (6)	0.64
II	4 (12)	2 (4)	0.23
III	11 (32)	3 (6)	<0.01
IV	18 (53)	39 (83)	<0.01
<b>Number of diseased vessels</b>			
1	22 (65)	31 (66)	1.00
2	8 (24)	11 (23)	1.00
3	4 (12)	5 (11)	1.00
Stent placed	31 (91)	44 (94)	0.69
Use of intra-aortic balloon pump	8 (24)	12 (26)	1.00
Creatinine pre-procedure, mg/dl	1.3 (0.9–1.6)	1.2 (1.0–1.7)	0.79
Hemoglobin pre-procedure, g/dl	11 ± 2	11 ± 3	0.33

Values are mean ± SD, n (%), or median (interquartile range).

PCI = percutaneous coronary intervention; STEMI = ST-segment elevation myocardial infarction.


patients with highest risk, whereas other investigators proposed that perhaps patients presenting in cardiogenic shock and cardiac arrest should be omitted from mandatory reporting (10). According to Massachusetts Data Analysis Center definitions, 22% (n = 18) of our patients who died met the compassionate use criteria; however, of these, only 4 had PCI-related mortality. Among the remaining 78% (n = 63), 52% (33 of 63) died of non-PCI-related mortality, highlighting the current limitation of excluding only those patients meeting the compassionate use definition.

Of the patients who died in our cohort, 58% (47 of 81 patients) presented with STEMI, cardiogenic shock, or cardiac arrest. This highlights the high-risk nature of many patients undergoing PCI who could potentially be considered emergent or salvage. Emergent procedures are not currently excluded from public reporting, and the current NCDR


(National Cardiovascular Data Registry) definition of salvage PCI would not incorporate many of these patients (21). In our cohort, the majority of these patients did not die of a PCI-related complication, perhaps indicating that such a presentation should not be a deterrent to PCI if otherwise indicated. Although these patients contribute significantly toward overall mortality, they are also individuals who are likely to benefit the most from PCI (30,31). Whereas some investigators have suggested excluding such patients from public reporting, using PCI- and non-PCI-related outcomes will allow public transparency yet avoid risk avoidance in these high-risk patients.

The main limitation of previous publications assessing PCI-associated mortality has been the sole reliance on death certificates that are well documented to be susceptible to serious biases and are often inaccurate (32–36). They often fail to capture other acute medical events during the course





of the hospital stay, which ultimately contributed to the death of the patient, thereby often mislabeling the etiology of death (37). Physicians often do not recall the cause of death and may confuse underlying cause of death with mechanism of death (20). In addition, such data may suffer from biases of ascertainment, whereby knowledge of a patient's prior medical conditions may affect physicians' interpretations of cause of death. For example, a patient's presentation with a myocardial infarction may bias a physician to label subsequent death as cardiac, even when it may not have been so. Similar to others' findings, death certificates had low sensitivity and specificity in our analysis. Therefore, it is reasonable to conclude that death certificates are an inaccurate representation of cause of death and hence should not be used for outcome reporting. There is no current consensus on the process of accurately determining



the cause of death, and questions on how the etiology of death should be established and who will ascertain it (hospital/healthcare professional performing the procedure or an independent committee) remain unanswered.

There are also other issues related to public reporting. Currently, there is no consensus on adjudicating death from anoxic brain injury after cardiac arrest into cardiac versus noncardiac death, especially if the patient underwent a successful PCI. Similarly, PCI-related complications such as bleeding or contrast-induced nephropathy may represent patient frailty rather than a direct complication of PCI. In our cohort, 4 patients had fatal bleeding; of these, only 1 was related to access site complication. Excluding the 3 patients with GI and intracranial bleeding would result in a lower incidence of PCI-related death. Furthermore, in many cases, a single cause of death cannot be identified due to ongoing multiorgan failure. Lastly, current public health reporting does not reward institutions that successfully treat high-risk patients who were refused previously nor report institutions that refuse PCI when it is indicated.

**Study limitations.** This is a single-center study and data from a tertiary referral hospital may not apply across other centers. Our study has a small sample size; however, our findings appear to be consistent with other studies that have assessed the accuracy of death certificates for determining cause of death. A conservative approach to adjudication of death as cardiac was made, and actual rates of cardiac death after PCI may be lower. We acknowledge the lack of standard definitions for several of the noncardiac causes of death across different registries, but we have made an attempt to use standard or common practice definitions wherever possible. All outside hospital cardiac arrests and deaths were considered as stent thrombosis and a PCI-related mortality; however, it is possible that a portion of these patients died from arrhythmias or other cardiac causes not related to PCI. Lastly, despite making every effort to be objective, unintentional errors in determining the exact cause of death may have occurred.

## Conclusions

Public reporting of PCI outcomes will allow transparency, help patients make informed decisions, and may be linked to reimbursements. However, in its current form, this may lead to risk avoidance and worse outcomes in high-risk patients. Classification of deaths into PCI-related and non-PCI-related may be a better marker of operator performance and quality of procedure. Death certificates have limitations and are not ideal for public health reporting.

## Acknowledgment

The authors thank Kathryn Brock for her editorial assistance.

**Reprint requests and correspondence:** Dr. Mehdi H. Shishehbor, Cleveland Clinic Foundation, 9500 Euclid Avenue, J3-5, Cleveland, Ohio 44195. E-mail: [shishem@ccf.org](mailto:shishem@ccf.org).

## REFERENCES

- Centers for Medicare and Medicaid Services. Proposals for improving quality of care during inpatient stays in acute care hospitals in the fiscal year 2011. Notice of proposed rule making. Available at: [http://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/AcuteInpatientPPS/downloads/FSQ09\\_IPLTCH11\\_NPRM041910.pdf](http://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/AcuteInpatientPPS/downloads/FSQ09_IPLTCH11_NPRM041910.pdf). Accessed April 1, 2012.
- U.S. Department of Health and Human Services. Hospital Compare. Available at: <http://www.naph.org/Main-Menu-Category/Publications/Quality/CMS-Quality-Data.aspx>. Accessed April 1, 2012.
- Werner RM, Asch DA. The unintended consequences of publicly reporting quality information. *JAMA* 2005;293:1239–44.
- Resnic FS, Welt FG. The public health hazards of risk avoidance associated with public reporting of risk-adjusted outcomes in coronary intervention. *J Am Coll Cardiol* 2009;53:825–30.
- Califf RM, Peterson ED. Public reporting of quality measures what are we trying to accomplish? *J Am Coll Cardiol* 2009;53:831–3.
- Joynt KE, Blumenthal DM, Orav EJ, Resnic FS, Jha AK. Association of public reporting for percutaneous coronary intervention with utilization and outcomes among Medicare beneficiaries with acute myocardial infarction. *JAMA* 2012;308:1460–8.
- Moscucci M. Public reporting of PCI outcomes and quality of care: one step forward and new questions raised. *JAMA* 2012;308:1478–9.
- Apolito RA, Greenberg MA, Menegus MA, et al. Impact of the New York State Cardiac Surgery and Percutaneous Coronary Intervention Reporting System on the management of patients with acute myocardial infarction complicated by cardiogenic shock. *Am Heart J* 2008;155:267–73.
- Adult percutaneous coronary intervention in the Commonwealth of Massachusetts J-D, 2005. Available at: <http://www.massdac.org/wp-content/uploads/CompUseCriteria.pdf>. Accessed April 1, 2012.
- Peterson ED, Dai D, DeLong ER, et al., for the NCDR Registry Participants. Contemporary mortality risk prediction for percutaneous coronary intervention: results from 588,398 procedures in the National Cardiovascular Data Registry. *J Am Coll Cardiol* 2010;55:1923–32.
- Ellis SG, Shishebor MH, Kapadia SR, et al. Enhanced prediction of mortality after percutaneous coronary intervention by consideration of general and neurological indicators. *J Am Coll Cardiol Intv* 2011;4:442–8.
- Resnic FS, Normand SL, Piemonte TC, et al. Improvement in mortality risk prediction after percutaneous coronary intervention through the addition of a “compassionate use” variable to the National Cardiovascular Data Registry CathPCI dataset: a study from the Massachusetts Angioplasty Registry. *J Am Coll Cardiol* 2011;57:904–11.
- Holmes DR, Selzer F, Johnston JM, et al., for the NHLBI Registry. Modeling and risk prediction in the current era of interventional cardiology: a report from the National Heart, Lung, and Blood Institute Dynamic Registry. *Circulation* 2003;107:1871–6.
- Singh M, Rihal CS, Lennon RJ, Garratt KN, Holmes DR Jr. A critical appraisal of current models of risk stratification for percutaneous coronary interventions. *Am Heart J* 2005;149:753–60.
- Rathore S, Grayson AD, Sastry S, et al., for the NWQIP Investigators. Circumstances and mode of in-hospital death following 9,914 consecutive patients undergoing percutaneous coronary interventions in the northwest of England. *J Invasive Cardiol* 2008;20:386–90.
- Busk M, Maeng M, Kristensen SD, et al., for the DANAMI-2 Investigators. Timing, causes, and predictors of death after three years’ follow-up in the Danish Multicenter Randomized Study of Fibrinolysis versus Primary Angioplasty in Acute Myocardial Infarction (DANAMI-2) trial. *Am J Cardiol* 2009;104:210–5.
- Stolker JM, Cohen DJ, Lindsey JB, Kennedy KF, Kleiman NS, Marso SP. Mode of death after contemporary percutaneous coronary intervention: a report from the Evaluation of Drug Eluting Stents and Ischemic Events registry. *Am Heart J* 2011;162:914–21.
- Cutlip DE, Windecker S, Mehran R, et al., for the Academic Research Consortium. Clinical end points in coronary stent trials: a case for standardized definitions. *Circulation* 2007;115:2344–51.
- Levy MM, Fink MP, Marshall JC, et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med* 2003;31:1250–6.
- Kircher T, Anderson RE. Cause of death: proper completion of the death certificate. *J Am Med Rec Assoc* 1987;58:47–51.
- National Cardiovascular Data Registry. NCDR CathPCI Registry v4. 4.Diagnostic catheterization and percutaneous coronary intervention registry. Available at: <https://www.ncdr.com/webncdr/cathpci/>. Accessed December 3, 2012.
- Malenka DJ, O’Rourke D, Miller MA, et al., for the Northern New England Cardiovascular Disease Study Group. Cause of in-hospital death in 12,232 consecutive patients undergoing percutaneous transluminal coronary angioplasty. *Am Heart J* 1999;137:632–8.
- King SB 3rd. A valuable public option. *J Am Coll Cardiol Intv* 2010;3:129–30.
- Levy RI, Jesse MJ, Mock MB. NHLBI position on percutaneous transluminal coronary angioplasty (PTCA). *Am J Cardiol* 1979;43:867.
- Omoigui NA, Miller DP, Brown KJ, et al. Outmigration for coronary bypass surgery in an era of public dissemination of clinical outcomes. *Circulation* 1996;93:27–33.
- Narins CR, Dozier AM, Ling FS, Zareba W. The influence of public reporting of outcome data on medical decision making by physicians. *Arch Intern Med* 2005;165:83–7.
- Matheny ME, Ohno-Machado L, Resnic FS. Discrimination and calibration of mortality risk prediction models in interventional cardiology. *J Biomed Inform* 2005;38:367–75.
- New York State Department of Health. Percutaneous coronary interventions (angioplasty) in New York State, 1995–1997 report. Available at: [http://www.health.ny.gov/statistics/diseases/cardiovascular/docs/pci\\_1995-1997.pdf](http://www.health.ny.gov/statistics/diseases/cardiovascular/docs/pci_1995-1997.pdf). Accessed April 1, 2012.
- New York State Department of Health. Percutaneous coronary interventions (angioplasty) in New York State 2001–2003 report. Available at: [http://www.health.ny.gov/statistics/diseases/cardiovascular/docs/pci\\_2001-2003.pdf](http://www.health.ny.gov/statistics/diseases/cardiovascular/docs/pci_2001-2003.pdf). Accessed April 1, 2012.
- Holmes DR Jr., Berger PB, Hochman JS, et al. Cardiogenic shock in patients with acute ischemic syndromes with and without ST-segment elevation. *Circulation* 1999;100:2067–73.
- Hochman JS, Sleeper LA, Webb JG, et al., for the SHOCK Investigators. Early revascularization in acute myocardial infarction complicated by cardiogenic shock. *N Engl J Med* 1999;341:625–34.
- Zumwalt RE, Ritter MR. Incorrect death certification: an invitation to obfuscation. *Postgrad Med* 1987;81:245–7. 250, 253–4.
- Lenfant C, Friedman L, Thom T. Fifty years of death certificates: the Framingham Heart Study. *Ann Intern Med* 1998;129:1066–7.
- Lloyd-Jones DM, Martin DO, Larson MG, Levy D. Accuracy of death certificates for coding coronary heart disease as the cause of death. *Ann Intern Med* 1998;129:1020–6.
- Harriss LR, Ajani AE, Hunt D, et al. Accuracy of national mortality codes in identifying adjudicated cardiovascular deaths. *Aust N Z J Public Health* 2011;35:466–76.
- Davignus ML, Liao Y, Greenland P, et al. Association of nonspecific minor ST-T abnormalities with cardiovascular mortality: the Chicago Western Electric Study. *JAMA* 1999;281:530–6.
- Ravakha K. Death certificates are not reliable: revivification of the autopsy. *South Med J* 2006;99:728–33.

**Key Words:** cause of death ■ coronary intervention ■ mortality ■ outcomes ■ public reporting ■ risk avoidance.